

2015

New distribution and host records for white coconut scale, *Parlagena bennetti* Williams (Hemiptera: Diaspididae)

Takumasa Kondo

Corporación Colombiana de Investigación Agropecuaria (CORPOICA), tkondo@corpoica.org.co

Penny Gullan

Australian National University, Canberra, penelope.gullan@anu.edu.au

Gillian W. Watson

Plant Pest Diagnostic Center, California Department of Food & Agriculture, gillian.watson@cdfa.ca.gov

Alex Enrique Bustillo Pardey

Programa de Plagas y Enfermedades Cenipalma, Colombia, abustillo@cenipalma.org

Luis Guillermo Montes

Entomología, Programa de Plagas y Enfermedades, Cenipalma, lmontes@cenipalma.org

Follow this and additional works at: <http://digitalcommons.unl.edu/insectamundi>

Kondo, Takumasa; Gullan, Penny; Watson, Gillian W.; Bustillo Pardey, Alex Enrique; and Montes, Luis Guillermo, "New distribution and host records for white coconut scale, *Parlagena bennetti* Williams (Hemiptera: Diaspididae)" (2015). *Insecta Mundi*. 927.
<http://digitalcommons.unl.edu/insectamundi/927>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0422

New distribution and host records for white coconut scale,
Parlagena bennetti Williams (Hemiptera: Diaspididae)

Takumasa Kondo

Corporación Colombiana de Investigación Agropecuaria (Corpoica)
Centro de Investigación Palmira, Calle 23, Carrera 37, Continuo al Penal
Palmira, Valle, Colombia

Penelope J. Gullan

Division of Evolution, Ecology & Genetics
Research School of Biology, The Australian National University
Canberra, A.C.T. 2601, Australia

Gillian W. Watson

Plant Pest Diagnostic Center, California Department of Food & Agriculture
3294 Meadowview Road
Sacramento, CA 95832-1437, U.S.A.

Alex Enrique Bustillo Pardey
Programa de Plagas y Enfermedades
Cenipalma, Colombia

Luis Guillermo Montes

Entomología, Programa de Plagas y Enfermedades, Cenipalma
Campo Experimental Palmar de la Vizcaína
Cenipalma-Zona Central, Colombia

Date of Issue: June 5, 2015

Takumasa Kondo, Penelope J. Gullan, Gillian W. Watson, Alex Enrique Bustillo Pardey, Luis Guillermo Montese
New distribution and host records for white coconut scale, *Parlagona bennetti* Williams (Hemiptera: Diaspididae)
Insecta Mundi 0422: 1–6

ZooBank Registered: urn:lsid:zoobank.org:pub:D7CCC1D9-9A76-4088-83EF-927E9FC9B7CF

Published in 2015 by

Center for Systematic Entomology, Inc.
P. O. Box 141874
Gainesville, FL 32614-1874 USA
<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Chief Editor: Paul E. Skelley, e-mail: insectamundi@gmail.com

Head Layout Editor: Eugenio H. Nearn

Editorial Board: J. H. Frank, M. J. Paulsen, Michael C. Thomas

Review Editors: Listed on the **Insecta Mundi** webpage

Manuscript Preparation Guidelines and **Submission Requirements** available on the **Insecta Mundi** webpage at: <http://centerforsystematicentomology.org/insectamundi/>

Printed copies (ISSN 0749-6737) annually deposited in libraries:

CSIRO, Canberra, ACT, Australia

Museu de Zoologia, São Paulo, Brazil

Agriculture and Agrifood Canada, Ottawa, ON, Canada

The Natural History Museum, London, UK

Muzeum i Instytut Zoologii PAN, Warsaw, Poland

National Taiwan University, Taipei, Taiwan

California Academy of Sciences, San Francisco, CA, USA

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

Field Museum of Natural History, Chicago, IL, USA

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Eugenio H. Nearn

New distribution and host records for white coconut scale, *Parlagena bennetti* Williams (Hemiptera: Diaspididae)

Takumasa Kondo

Corporación Colombiana de Investigación Agropecuaria (Corpoica)
Centro de Investigación Palmira, Calle 23, Carrera 37, Continuo al Penal
Palmira, Valle, Colombia
tkondo@corpoica.org.co

Penelope J. Gullan

Division of Evolution, Ecology & Genetics
Research School of Biology, The Australian National University
Canberra, A.C.T. 2601, Australia
penelope.gullan@anu.edu.au

Gillian W. Watson

Plant Pest Diagnostic Center, California Department of Food & Agriculture
3294 Meadowview Road
Sacramento, CA 95832-1437, U.S.A.
gillian.watson@cdfa.ca.gov

Alex Enrique Bustillo Pardey

Programa de Plagas y Enfermedades
Cenipalma, Colombia
abustillo@cenipalma.org

Luis Guillermo Montes

Entomología, Programa de Plagas y Enfermedades, Cenipalma
Campo Experimental Palmar de la Vizcaína
Cenipalma-Zona Central, Colombia
lmontes@cenipalma.org

Abstract. The white coconut scale, *Parlagena bennetti* Williams, 1969 (Hemiptera: Diaspididae) is reported for the first time on New Zealand flax, *Phormium tenax* J.R. Forst. & G. Forst. (Xanthorrhoeaceae), coconut, *Cocos nucifera* L., Manila palm, *Veitchia merrillii* (Becc.) H.E. Moore, oil palm, *Elaeis guineensis* Jacq. and an oil palm hybrid OxG: *E. oleifera* x *E. guineensis* (Arecaceae) in continental Colombia, and on coconut from Venezuela. Previously, *P. bennetti* was known only on the Caribbean islands of Trinidad and San Andres, causing serious damage to coconut, its only known host until now. A brief characterization of *P. bennetti* and an updated list of 23 species of scale insects of the family Diaspididae reported worldwide on *Elaeis* spp. is provided.

Key words. Armored scale, Colombia, *Elaeis guineensis*, insect pest, oil palm, species list, Venezuela.

Resumen. Se reporta por primera vez la presencia de la escama blanca del cocotero *Parlagena bennetti* Williams, 1969 (Hemiptera: Diaspididae) sobre lino de Nueva Zelanda, *Phormium tenax* J.R. Forst. & G. Forst. (Xanthorrhoeaceae), palma de coco *Cocos nucifera* L., palma Manila *Veitchia merrillii* (Becc.) H.E. Moore, palma de aceite *Elaeis guineensis* Jacq. y un híbrido de palma de aceite OxG: *E. oleifera* x *E. guineensis* (Arecaceae) en Colombia continental y sobre cocotero en Venezuela. Hasta ahora *P. bennetti* solo se conocía en la isla de Trinidad y en San Andrés, en el Caribe, causando graves daños al cocotero, el único hospedero conocido hasta ahora. Se provee una breve caracterización de *P. bennetti*, y un listado actualizado de 23 insectos escama de la familia Diaspididae reportadas sobre *Elaeis* spp. en el mundo.

Palabras Clave. Escama de armadura, Colombia, *Elaeis guineensis*, insecto plaga, listado de especies, palma de aceite, Venezuela.

Introduction

Scale insects (Hemiptera: Sternorrhyncha: Coccoidea) are small sap-sucking insects with soft bodies, with about 8000 described species in more than 30 families worldwide (Ouvrard et al. 2013). Recently, the first author received samples of an armored scale causing damage to oil palm in the State of Santander, in northeastern Colombia. The scale insect samples were slide-mounted by the second author and identified as *Parlagena bennetti* Williams through the use of the original description by Williams (1969), and assistance from the third author. Later, the first author inspected palm species and other monocots planted in the grounds of the Colombian Corporation for Agricultural Research, Palmira Research Station and neighboring areas in the city of Palmira, State of Valle del Cauca, and found the same species on leaves and fruit of coconut, Manila palm, New Zealand flax and on leaves of a hybrid oil palm, OxG: *Elaeis oleifera* x *E. guineensis*. Slide-mounted specimens from Venezuela on coconut were also found in the California State Collection of Arthropods during this study. Hitherto, *P. bennetti* was recorded in the literature only on coconut from the Caribbean islands of Trinidad (Williams 1969) and San Andres (Mosquera 1976).

Heavy infestations of armored scale insects on palm fronds often cause leaf chlorosis, premature senescence, patches of necrotic leaf tissue, and premature leaf drop (Beardsley and González 1975). According to Williams (1969), Dr. F. Bennett described an infestation by *P. bennetti* on coconut in Trinidad as being associated with presence of necrotic areas on the leaves, with heavily attacked palms having an unthrifty appearance that was probably due to plant damage.

In mainland Colombia ten species of armored scale insects (Hemiptera: Diaspididae) have been reported previously on oil palm (*E. guineensis*) namely, *Aonidiella orientalis* (Newstead), *Aspidiotus destructor* Signoret, *A. excisus* Green, *Diaspis boisduvalii* Signoret, *Hemiberlesia lataniae* (Signoret), *Hemiberlesia* sp. possibly *H. palmae* (Cockerell), *Ischnaspis longirostris* (Signoret), *Pinnaspis aspidistrae* (Signoret), *P. strachani* (Cooley), and *Selenaspis articulatus* (Morgan) (Posada-Ochoa 1989).

The purpose of this paper is to report for the first time the presence of *P. bennetti* in continental Colombia and Venezuela, and also record New Zealand flax, *Phormium tenax* J.R. Forst. & G. Forst. (Xanthorrhoeaceae), Manila palm, *Veitchia merrillii* (Becc.) H.E. Moore, oil palm, *Elaeis guineensis* L. and an oil palm hybrid OxG, *E. oleifera* and *E. guineensis* (Arecaceae) as new hosts for this armored scale. An updated list of armored scale insects recorded on oil palm is provided.

Materials and Methods

Adult female specimens were slide-mounted following the procedure described by Williams and Granara de Willink (1992) except that 100% isopropyl alcohol (= 2-propanol) was used for dehydration and xylene was used instead of clove oil. Morphological terms for armored scales follow Williams and Watson (1988) and Miller and Davidson (2005). Photographs were taken *in situ* with a Canon Ixy 630 digital camera and close-ups of the insect were taken using the same camera with the aid of a stereomicroscope Nikon SMZ 745T. Photomicrographs of an adult female were made using a Nikon Digital




Figure 1. The white coconut scale, *Parlagena bennetti*. **A).** Infesting coconut fruits. **B).** Close-up of insects infesting fruit. **C).** Typical infestation on leaves. Photographs by T. Kondo, Palmira, Colombia.

Sight DS-5M camera, NIS-Elements F 3.0 software, and a Nikon Eclipse 80i compound microscope. Auto-montage of a photomicrograph series was done using Combine ZP software. The images were processed using the computer program Adobe Photoshop® CS 5.1.

The material studied is deposited in the Museo de Entomología, Corporación Colombiana de Investigación Agropecuaria, Centro de Investigación Palmira, Palmira, Valle del Cauca, Colombia (MECP), the California State Collection of Arthropods at the Plant Pest Diagnostics Center, California Department of Food & Agriculture, 3294 Meadowview Road, Sacramento, California, U.S.A. (CSCA) and the Museo Entomológico, Facultad de Agronomía, Universidad Nacional de Colombia, Sede Bogotá, Bogotá, Cundinamarca, Colombia (UNAB).

Taxonomy

Parlagena bennetti Williams, 1969.

Parlatoria bennetti (Williams), Davies and Boratynski 1979. [This change of combination is not accepted.]

Characterization of adult female. The following characterization of *P. bennetti* is based on the original description by Williams (1969).

Insects in life (Fig. 1A–1C). Scale cover of adult female white, about 1.00–1.25 mm in diameter, dome-shaped, with a distinctly downy texture. Exuviae pale yellow, situated at edge of scale, often at right angles to scale and leaf surface. Adult female small, 0.6 mm long and dark brown, completely surrounded by downy wax.

Slide-mounted specimens (Fig. 2A, 2B). Adult female about 0.6 mm in length, slightly longer than wide, turbinate. Body membranous except for pygidium. Spiracles lacking associated pores. Pygidium rounded. Median lobes small, short and rounded, about as long as wide, with a faint suggestion of a notch on outer margin; separated by a space greater than width of a lobe. Second lobes bilobed, varying in shape but always short and rounded. Third lobes represented by small projections from margin. Plates indistinct and very fragile, scarcely longer than a median lobe, with a pair between median lobes and one between each median and second lobe; plates without associated microducts.

Dorsal macroducts numerous and scattered; each orifice oval, surrounded by a distinct sclerotized rim, set almost at right angles to body margin. Each duct two-barred, with distal or innermost bar thickest and space between bars sclerotized. Marginal ducts near lobes situated a short distance from margin, with one or two between median lobes. Other ducts on pygidium scattered, becoming progressively smaller as far forward as second abdominal segment. Anal ring elongate, widest at anterior end, situated about one-third length of pygidium from apex. Perivulvar pores in five groups; median group with 1–4 pores, each anterior lateral group with 5–8 pores and each posterior lateral group with 2–5




Figure 2. A). Auto-montage photomicrograph of an adult female of *Parlagena bennetti* from Venezuela, Lara, Barquisimeto. Photomicrograph by G.W. Watson, California Department of Food & Agriculture. B). Illustration of body and details of the pygidium (after Williams 1969).

Table 1. Updated list of armored scale insects (Hemiptera: Diaspididae) recorded on oil palm, *Elaeis* spp. (Arecaceae) in the world (adapted from Howard 2001 and Ben-Dov et al. 2015).

Diaspidid species	Literature records on <i>Elaeis</i>
<i>Aonidiella gracilis</i> (Balachowsky)	Balachowsky (1956)
<i>Aonidiella rex</i> Balachowsky	Balachowsky (1956), Howard (2001)
<i>Aspidiotus destructor</i> Signoret	Castel-Branco (1963), Almeida (1973), Williams and Watson (1988), Howard (2001)
<i>Aspidiotus elaeidis</i> Marchal	Marchal (1909), Sanders (1909), Leonardi (1914), Balachowsky (1956), Howard (2001)
<i>Aspidiotus nerii</i> Bouché	Howard (2001)
<i>Aspidiotus remaudierei</i> Balachowsky	Balachowsky (1956)
<i>Chrysomphalus dictyospermi</i> (Morgan)	Almeida (1973), Williams and Watson (1988)
<i>Diaspis boisduvalii</i> Signoret	Miller and Davidson (2005)
<i>Diaspis elaeidis</i> Munting	Munting (1969)
<i>Hemiberlesia lataniae</i> (Signoret)	Balachowsky (1956), Munting (1969), Almeida (1971, 1973), Howard (2001)
<i>Hemiberlesia palmae</i> (Cockerell)	Almeida (1973), Howard (2001)
<i>Ischnaspis longirostris</i> (Signoret)	Mamet (1943), Ben-Dov (1974), Miller and Davidson (2005)
<i>Leucaspis fulchironiae</i> (Boisduval)	Boisduval (1867)
<i>Lindingaspis musae</i> (Laing)	Balachowsky (1958), Howard (2001)
<i>Lindingaspis opima</i> (Silvestri)	Almeida (1973)
<i>Parlagentia bennetti</i> Williams	Present study
<i>Parlatoria proteus</i> (Curtis)	Barber (1910)
<i>Pinnaspis aspidistrae</i> (Signoret)	Williams and Watson (1988)
<i>Pinnaspis strachani</i> (Cooley)	Cockerell (1902), Ferris and Rao (1947), Williams and Watson (1988)
<i>Pseudaonidia trilobitiformis</i> (Green)	Almeida (1973)
<i>Pseudaulacaspis cockerelli</i> (Cooley)	Chen (1983)
<i>Rolaspis chaetachmae</i> (Brain)	Munting (1969)
<i>Selenaspis articulatus</i> (Morgan)	Mamet (1958), Borchsenius (1966), Almeida (1973), Howard (2001)

pores. Microducts not numerous. Vulva situated almost in center of pygidium. Sclerotized duct tubercles absent.

Material studied. Colombia: Santander: Barrancabermeja, 6.ii.2015, coll. L. Montes, ex foliage of *Elaeis guineensis* Jacq., 3 slides, 14 specimens (MECP). Valle del Cauca: Palmira, Hotel las Victorias, Carrera 32, via a Candelaria, contiguo a la Universidad Nacional de Colombia, sede Palmira, 03°30'43.0"N, 76°18'46.0"W, 1020 m a.s.l., 17.iv.2015, coll. T. Kondo, ex leaves of *Phormium tenax* J.R. Forst. & G. Forst, 1 slide, 4 specimens (MECP); Palmira, near Centro Comercial Llanogrande, 03°31'44.6"N, 76°19'08.3"W, 1009 m a.s.l., 2.iv.2015, coll. T. Kondo, ex leaves of *Veitchia merrillii* (Becc.) H.E. Moore, 1 slide, 5 specimens (UNAB); Palmira, Corporación Colombiana de Investigación Agropecuaria (Corpoica), Centro de Investigación Palmira, Banco de germoplasma de frutales tropicales, 03°30'49.5"N, 76°19'43.5"W, 1001 m a.s.l., coll. 2.iv.2015, coll. T. Kondo, ex fruit of *Cocos nucifera* L., 3 slides, 15 specimens (UNAB); Palmira, Corporación Colombiana de Investigación Agropecuaria (Corpoica), Centro de Investigación Palmira, 03°30'48.3"N, 76°19'03.9"W, 1010 m a.s.l., coll. 3.iv.2015, coll. T. Kondo, ex leaves of OxG hybrid oil palm, *E. oleifera* x *E. guineensis*, 1 slide, 5 specimens (UNAB). Venezuela: Lara: Barquisimeto, v.2006, coll. R. Urtiaga, ex *C. nucifera*, 3 slides, 15 specimens (CSCA). Trinidad: Manzanilla Beach, 14.xi.1995, coll. M.J.W. Cock, ex leaves of *C. nucifera*, 1 slide, 3 specimens (CSCA).

Distribution. Colombia: San Andrés y Providencia, Santander, Valle del Cauca. **Trinidad and Tobago:** Trinidad. **Venezuela:** Lara.

Host plants. Arecaceae: *Cocos nucifera*, *Elaeis guineensis*, hybrid oil palm, *E. oleifera* x *E. guineensis*, *Veitchia merrillii*. **Xanthorrhoeaceae:** *Phormium tenax*.

Remarks. *Parlagentia bennetti* may infest fruit, covering much of the fruit surface if the infestation is heavy (Fig. 1A, B); however, usually the scales are found on the underside of the leaves (Fig. 1C) although sometimes they occur on both surfaces. Williams (1969) described the adult females as dark and heavily sclerotized, however, it should be noted that young adult females are grayish white in color and membranous. The scale cover is somewhat shell-like (Fig. 1B). The scale of the adult male is about 1.0 mm long, elongate, with sides subparallel, and of the same downy consistency as female scale (Williams 1969) (Fig. 1B). The discovery of *P. bennetti* on a non-palm species, i.e., *Phormium tenax* (Xanthorrhoeaceae), indicates that the scale insect may be able to attack other species of monocots. No natural enemies have been reported in the literature.

Armored scales recorded on oil palm, *Elaeis* spp. There were 22 species of armored scales recorded previously on oil palms, *Elaeis* spp., worldwide. With the addition of *P. bennetti*, this tally is increased to 23 species (Table 1).

Acknowledgments

Many thanks to Mr. Silvio Bastidas (Corpoica, Palmira Research Station) for the identification of the oil palm hybrid. The authors thank Dr. Lucia Claps (Universidad Nacional de Tucumán, Argentina) and Dr. Vera Wolff (FEPAGRO - Fundação Estadual de Pesquisa Agropecuária, Porto Alegre, Rio Grande do Sul, Brazil) for reviewing the manuscript. We are grateful to Mr. Alejandro Caballero for slide-mounting the material deposited at UNAB.

Literature Cited

- Almeida, D. M. de 1971.** Diaspididae (Homoptera - Coccoidea) of Mozambique, with the description of a new species. *Novos Taxa Entomológicos* 90: 3–17.
- Almeida, D. M. de 1973.** Coccoidea de Angola. 1- Revisão das espécies conhecidas. *Boletim do Instituto de Investigação Científica de Angola* 10: 1–23.
- Balachowsky, A. S. 1956.** Les cochenilles du continent Africain Noir. V. 1 - Aspidiotini (1ère partie). *Annales du Musée Royal du Congo Belge (Sciences Zoologiques)* 3: 1–142.
- Balachowsky, A. S. 1958.** Les cochenilles du continent Africain Noir. v. 2 Aspidiotini (2me partie), Odonaspidini and Parlatorini. *Annales du Musée Royal du Congo Belge (Sciences Zoologiques)* 4: 149–356.
- Barber, T. C. 1910.** The Coccidae of Audubon Park, New Orleans. *Journal of Economic Entomology* 3: 420–425.
- Beardsley, J. W. Jr., and R. H. González. 1975.** The biology and ecology of armored scales. *Annual Review of Entomology* 20: 47–73.
- Ben-Dov, Y. 1974.** A revision of *Ischnaspis* Douglas with a description of a new allied genus (Homoptera: Diaspididae). *Journal of Entomology (B)* 43: 19–32.
- Ben-Dov, Y., D. R. Miller, and G. A. P. Gibson. 2015.** ScaleNet: a database of the scale insects of the world. (Available at ~ <http://www.sel.barc.usda.gov/scalenet/scalenet.html>). Last accessed February, 2015.)
- Boisduval, J. B. A. 1867.** *Essai sur l'entomologie horticole*. Donnad; Paris, Fr. 648 p.
- Borchsenius, N. S. 1966.** A catalogue of the armoured scale insects (Diaspidoidea) of the world. Nauka, Moscow and Leningrad. 449 p.
- Castel-Branco, A. J. F. 1963.** *Aspidiotus destructor* Sign. and other scales that occur on coconut trees (*Cocos nucifera* L.) and African oil palm (*Elaeis guineensis* Jacq.) on the islands of S. Tome and Principe. *Memorias da Junta de Investigações do Ultramar (2a series)* 43: 131–175.

- Chen, F. G. 1983.** The Chionaspidini (Diaspididae, Coccoidea, Homoptera) from China. Science & Technology Publishing House; Sichuan Province, China. 175 p.
- Cockerell, T. D. A. 1902.** Note sur le genre *Hemichionaspis* [Coccid.] et description d'une espèce nouvelle. Bulletin de la Société Entomologique de France 1902: 81–82.
- Davies, R. G., and K. L. Boratynski. 1979.** Character selection in relation to the numerical taxonomy of some male Diaspididae (Homoptera: Coccoidea). Biological Journal of the Linnaean Society 12: 95–165.
- Ferris, G. F., and V. P. Rao. 1947.** The genus *Pinnaspis* Cockerell (Homoptera: Coccoidea: Diaspididae). (Contribution No. 54). Microentomology 12: 25–58.
- Howard, F. W. 2001.** Hemiptera: Sternorrhyncha. p. 161–227. In: F. W. Howard, D. Moore, R. M. Giblin-Davis, and R. G. Abad. Insects on palms. Chapter 3. Sap-feeders on palms. CABI Publishing; Wallingford, UK, 400 p.
- Leonardi, G. 1914.** Contributo alla conoscenza delle coccinglie dell'Africa occidentale e meridionale. Bollettino del Laboratorio di Zoologia Generale e Agraria della R. Scuola Superior Agricoltura. Portici 8: 187–224.
- Mamet, J. R. 1943.** A revised list of the Coccoidea of the islands of the western Indian Ocean, south of the equator. Mauritius Institute Bulletin 2: 137–170.
- Mamet, J. R. 1958.** The *Selenaspidus* complex (Homoptera, Coccoidea). Annales du Musée Royal du Congo Belge, Zoologiques, Miscellanea Zoologica 4: 359–429.
- Marchal, P. 1909.** Cochenilles nouvelles de l'Afrique occidentale. Bulletin de la Société Zoologique de France 34: 68–69.
- Miller, D. R., and J. A. Davidson. 2005.** Armored scale insect pests of trees and shrubs. Cornell University Press; Ithaca, NY. 442 p.
- Mosquera, P. F. 1976.** Escamas protegidas más frecuentes en Colombia. Boletín Técnico, Ministerio de Agrícola Instituto Colombiano Agropecuario, División de Sanidad Vegetal 38: 1–103.
- Munting, J. 1969.** Observations on some armoured scale insects (Homoptera: Coccoidea: Diaspididae) from south west Africa and neighbouring territories. Cimbebasia 1 (ser. A): 115–161.
- Ouvrard, D., T. Kondo, and P. J. Gullan. 2013.** Scale insects: major pests and management. Encyclopedia of Pest Management, pp. 1–4 [published online: 1 March 2013. Taylor and Francis; New York, NY. DOI: 10.1081/E-EPM-120046899].
- Posada-Ochoa, L. 1989.** Lista de insectos dañinos y otras plagas en Colombia. Cuarta Edición, Boletín Técnico No.43. Instituto Colombiano Agropecuario. 662 p.
- Sanders, J. G. 1909.** Catalogue of recently described Coccidae - II. United States Department of Agriculture, Bureau of Entomology, Technical Series 16: 33–60.
- Williams, D. J. 1969.** A new species of *Parlagena* McKenzie attacking coconut trees in Trinidad (Hom., Coccoidea, Diaspididae). Bulletin of Entomological Research 59: 97–99.
- Williams, D. J., and M. C. Granara de Willink. 1992.** Mealybugs of Central and South America. CAB International; Wallingford, U.K. 635 p.
- Williams, D. J., and G. W. Watson. 1988.** The scale Insects of the Tropical South Pacific Region. Part 1. The armoured scales (Diaspididae). CAB International; Wallingford, U.K. 290 p.

Received April 9, 2015; Accepted May 21, 2015.

Review Editor Aline Barcellos.